arXiv:1304.8116v1 [astro-ph.GA] 30 Apr 2013

Pulsar scintillation patterns and strangelets

M. Ángeles Pérez-García[2] , Joseph Silk[1] and Ue-Li Pen[3]
[1] *Institut d'Astrophysique de Paris, F-75014 Paris, France,*
[2] *Department of Fundamental Physics and IUFFyM,*
Universidad de Salamanca, Plaza de la Merced s/n 37008 Salamanca,
[3] *Canadian Institute for Theoretical Astrophysics, University of Toronto, 0N M5S 3H8, Canada*
(Dated: May 1, 2013)

We propose that interstellar extreme scattering events, usually observed as pulsar scintillations, may be caused by a coherent agent rather than the usually assumed turbulence of H_2 clouds. We find that the penetration of a flux of ionizing, positively charged strangelets or quark nuggets into a dense interstellar hydrogen cloud may produce ionization trails. Enhanced electron densities would then form and constitute a lens-like scattering screen for radio pulsars and quasars.

A variety of scintillation phenomena observed from pulsars and quasars require interstellar scattering screens that contain compact regions of high electron density. These include quasar Extreme Scattering Events (ESE) [1, 2], pulsar parabolic arcs [3, 4] and Galactic Center scattering of OH maser sources [5, 6]. Many of these phenomena, in particular the ESEs, require enhanced electron density regions of A.U. ($\sim 1.5\,10^{13}$ cm) size. The overpressure $P/k_B \approx 10^6 - 10^8\,\mathrm{K\,cm^{-3}}$, as estimated from typical temperatures $T \sim 10^4\,\mathrm{K}$ and particle densities $n \sim 10^2 - 10^4\,\mathrm{cm^{-3}}$, is difficult to explain in any conventional scattering screens embedded in dense molecular H_2 clouds. The only plausible environment where such pressures might be attained would be in the dense cloud cores (cf. the compact ionized cloud model developed by Walker [7, 8]). The required source properties in this latter model require a significant mass in extremely dense cold gas clumps to source the ionized gas. The stability of such cold clumps is questionable, although exotic models have been proposed [9].

There is some direct evidence that the ionized clouds are highly elongated [10]. It is an already well-known fact that plasma lenses result from electron (over-) under-densities. Electron over-densities result in a faster phase velocity, corresponding to a concave (divergent) optical lens, while under-densities are associated with a convergent lens [11]. Quantitative lens models for the ESEs [7] estimate column densities $\rho_e \sim 10^{15}\,\mathrm{cm^{-2}}$.

Here we are interested in estimating the effect of the enhancement of the electron column density on A.U. scales as a result of the formation of ionized trails in HI and H_2 clouds. We propose a mechanism that provides an alternative to postulating the existence of controversial clumps of dense molecular hydrogen, and naturally generates pervasive ionized trails in dense interstellar cloud cores. Our model invokes strangelets [12] (also known as *nuclearites*), finite droplets of quark matter with non-zero strangeness fraction and slightly charged. They are currently being searched on earth [13] for as final products in heavy ion collisions with the ALICE experiment at the LHC, in the CDMSII under the form of light ionizing particles or in the AMS-02 mission. It is expected that these quark droplets can be naturally generated by a series of different astrophysical events where a nucleon-quark deconfinement transition may take place, e.g. neutron star (NS) collisions, NS or black hole combined binary mergers or in NS to quark star (QS) conversions. This latter process may be induced by internal heating due to dark matter (DM) annihilations [14] (under the assumption of a Majorana DM particle candidate) and even leave observable traces in the pulsar distribution [15] or in the emission of very short gamma ray bursts (GRBs) with typical time scales $T_{90} \lesssim 0.1$ s [16]. Due to the large gravitational and nuclear binding energies released in the transitioning process, a mass ejection episode is expected, possibly seeding the interstellar medium with a fraction of strangeness-carrying lumps of matter formed during the phase of nucleon deconfinement. Energetics for short GRBs isotropic equivalent photon emission show that the measured value $E_{\gamma\, iso} \sim 10^{48} - 10^{52}$ erg is compatible with relativistic mass ejecta $M_{\rm ej} \lesssim 10^{-4} M_\odot$ able to consistently produce observable gamma rays. It has been actually proposed that quark matter droplets might partially populate cosmic ray (CR) primaries (see e.g. [17–19]).

In this context, let us consider a volume of interstellar space that initially contains a ~ 30 pc cloud radius of H_2 (molecules) and HI (neutral) atoms. Typical ionization reactions are of the form, $\mathrm{X} + \mathrm{H_2} \rightarrow \mathrm{X} + \mathrm{H_2^+} + \mathrm{e^-}$, or $\mathrm{X} + \mathrm{HI} \rightarrow \mathrm{X} + \mathrm{H^+} + \mathrm{e^-}$, where X is the incoming charged strangelet (assumed to have unit charge in the previous reaction, but more generally with charge Ze). The energy needed to ionize a hydrogen atom (molecule), initially in the ground state is I(HI) = 13.6 eV (I(H_2) = 15.6 eV). In astrophysical CGS units we use the conversion factor $1\,\mathrm{eV} = 1.62 \times 10^{-12}$ erg. The dimensions of the cloud can be estimated assuming a radius for the cloud core $\mathrm{R_C} \sim 10\,\mathrm{pc} \sim 3 \times 10^{19}$ cm, then the core volume is $\mathrm{V_C} \approx 4\pi \mathrm{R_C^3}/3 \simeq 1.1 \times 10^{59}$ cm^3. Since the atomic number density in the cloud core is $n_0 \sim 10^4$ cm^{-3}, the number of particle species in the original neutral cloud (HI, protons, electrons) is accordingly given by $n_0\, V_C \simeq 1.1 \times 10^{63}$. In general, other ionizing agents, such as CRs can cause ionization of H_2 or HI with an



FIG. 1. Physical scenario of ionization in the H_2 cloud due to an emitting source of strangelets in the line of sight to the observer.

ionization rate $\xi^{HI} \sim 10^{-15}\,\mathrm{s}^{-1}$ [20] or $\xi^{H_2} \sim 10^{-17}\,\mathrm{s}^{-1}$ [21]. A more exotic flux of potentially ionizing electrically charged strangelets through the cloud will depend, on the one hand, on the time and spatial distribution of their astrophysical sources (besides a possible primordial background) and, on the other hand, on their peculiar nature.

An estimate of the incoming flux starts with the strangelet formation rate associated with the different astrophysical sources involved. We will consider a simplified model with a unique A baryonic number lump of mass $m^A_{\mathrm{STR}} < A m_{\mathrm{N}}$, where m_{N} is the nucleon mass. The strangelet number production rate in the astrophysical ith-process will be given by $dN^i_{\mathrm{A}}/dt = \eta_i \dot{M}_{\mathrm{ej,i}}/m^{\mathrm{A}}_{\mathrm{STR}}$. $\dot{M}_{\mathrm{ej,i}}$ is the mass rate ejected in the ith-process and η_i is the efficiency of strangelet ejection for the ith-process involved. For example, for NS collisions it is expected that $M_{\mathrm{ej}} \sim (10^{-5} - 10^{-2})M_\odot$ while in a NS merger the mass ejection is $M_{\mathrm{ej}} \sim (5\times 10^{-4} - 7\times 10^{-3})M_\odot$ for equal-mass binaries with total mass $m = 2.7 M_\odot$ [22]. In a NS to QS transition capable of emitting a GRB, it is expected that $M_{\mathrm{ej}} \lesssim 10^{-4} M_\odot$. According to [16] the rate of these transitions is $R_{\mathrm{NS-QS}} \simeq (8\times 10^{-4} - 3\times 10^{-3}) R_{\mathrm{SNtII}}$ being $R_{\mathrm{SNtII}} \simeq 10^{-2}\ \mathrm{yr}^{-1}$ the rate of type II supernovae in our galaxy.

Having enumerated the possible processes that may constitute a source of the strangelet flux, for the rest of this work we will consider a *generic source* where the deconfinement transition can take place with galactic appearance rate $R = \bar{R} \simeq 10^{-5}\ \mathrm{yr}^{-1}$ and ejected mass $M_{\mathrm{ej}} = \bar{M}_{\mathrm{ej}} \simeq 10^{-5} M_\odot$. The ejected mass rate is then $\dot{M}_{\mathrm{ej}} = R\, M_{\mathrm{ej}} \simeq 10^{-10} M_\odot\,\mathrm{yr}^{-1}$. Since the efficiency of strangelet production depends on the so far unknown details of the engine model, we will consider that only a small fraction $\eta \sim 10^{-2}$ is ejected under exotic form [16]. Nevertheless, strangelets should be emitted with A-values larger than a critical stability value [23], $A > A_{\mathrm{min}} \simeq 10^1 - 10^2$ so that they can possibly decay to the lightest energetically stable A_{min}-species.

The strangelet number production rate at the source is

$$\frac{dN_A}{dt} = 2\times 10^{45}\left(\frac{\eta}{0.01}\right)\left(\frac{\dot{M}_{\mathrm{ej}}}{10^{-10} M_\odot\,\mathrm{yr}^{-1}}\right) f_S(Z,\beta) A^{-1}\,\mathrm{yr}^{-1}. \tag{1}$$

It is expected that a possible emission distribution function at the source $f_S(Z,\beta)$ can modulate this rate. We will not consider this refinement here, and in what follows we will assume $f_S(Z,\beta) = 1$. From such a source, the corresponding unscreened flux measured by an observer at distance $d_{\mathrm{SO}} \sim 600$ pc is given by

$$F = \frac{dN_A}{dt}\frac{1}{4\pi d_{SO}^2} \sim 1.6\times 10^{-6} A^{-1}\,\mathrm{cm}^{-2}\mathrm{sr}^{-1}\mathrm{s}^{-1}, \tag{2}$$

for $A \gtrsim A_{\mathrm{min}}$. Note that for strangelets being currently searched for in neutrino telescopes on earth, there is a lower limit $A \gtrsim 10^{13}$ GeV/c^2. Our estimates yield in that case $F \lesssim 1.6\times 10^{-19}\,\mathrm{cm}^{-2}\mathrm{sr}^{-1}\mathrm{s}^{-1}$. Unfortunately, this is not presently within the reach of ANTARES, who report[24] a testing capability of 90% C. L. flux limit of $F_{\mathrm{ANTARES}} \sim 2\times 10^{-14}\,\mathrm{cm}^{-2}\mathrm{sr}^{-1}\mathrm{s}^{-1}$ for $A \gtrsim 10^{13}$, or IceCube-22, who report $F_{\mathrm{IC22}} \sim 10^{-18}\,\mathrm{cm}^{-2}\mathrm{sr}^{-1}\mathrm{s}^{-1}$ for $A \gtrsim 10^{17}$ nuclearites [25].

Once the nuclearites are produced it may happen that their trajectories intersect with a H_2 cloud along the line of sight to the observer, as depicted in Fig. 1. In this scenario one can estimate N^{eff}_A, the number of ionizing $A-$sized strangelets traversing the cloud in its lifetime, $\tau_c \sim 10^7$ yr, as $N^{\mathrm{eff}}_A = \eta_\Omega \frac{\pi R_C^2}{4\pi d_{SC}^2}\int_0^{\tau_c}\frac{dN_A}{dt}dt$. Using typical values [26] for the distances in this expression, we obtain

$$N^{\mathrm{eff}}_A \simeq 1.25\,10^{45} A^{-1}\left(\frac{\eta_\Omega}{0.01}\right)\left(\frac{R_C}{1\,\mathrm{pc}}\right)^2\left(\frac{200\,\mathrm{pc}}{d_{\mathrm{SC}}}\right)^2\left(\frac{\tau_c}{10^7\,\mathrm{yr}}\right), \tag{3}$$

where d_{SC} is the distance from the source to the cloud. Since beaming is observed in some very short GRBs (expected in this scenario [16]) with opening angle $\theta_{\mathrm{j}} \sim 1^\circ - 30^\circ$, an efficiency factor $\eta_\Omega \simeq \frac{\theta_{\mathrm{j}}^2}{2}$ has been introduced.

The peculiar nature of the quark droplets is highly uncertain but it is usually assumed that their charge is small and distributed positively on the surface [17]. For ordinary strangelets $Z = 0.3A^{1/3}$ [27] while for CFL (color-flavour-locked) strangelets $Z \simeq 0.3A^{2/3}$. Even smaller charge-to-mass ratios are energetically possible for intermediate masses, $A \sim 10^2 - 10^{18}$, assuming a strong coupling $\alpha_S = 0.9$ [28]. For example, for $A \simeq 10^9$, $Z/A \simeq 10^{-4}$ while larger strangelets $A \simeq 10^{18}$ have $Z/A \simeq 10^{-7} - 10^{-3}$ and even $Z/A < 0$. In this work we constrain strangelets to have $Z \geq 1$.

As the slightly charged and heavy strangelet traverses the hydrogen cloud in nearly straight lines, it continuously loses kinetic energy. Typically, the ejection energy at the source allows these particles to achieve a Lorentz factor bounded by a saturation value, $\gamma \lesssim \gamma_{\mathrm{sat}}$ with



FIG. 2. Logarithm of the kinetic energy loss rate in the cloud as a function of the logarithm of the incoming ($A = 10^3$) strangelet kinetic energy per baryon number. Several charge states are shown and a CR proton case with $Z = 1$ is depicted for comparison.



FIG. 3. Logarithm of the strangelet range in the cloud as a function of the logarithm of the incoming kinetic energy per baryon number. We consider a CR proton $Z = 1$, ordinary and CFL strangelets with $A = 10^2$ and $A = 10^6$ respectively, and larger strangelets with $A = 10^{13}$ and $A = 10^{18}$ with $Z/A = 10^{-7}$.

$\gamma_{\rm sat} \sim 20 - 1000$ [16]. Strangelets of net effective charge Z_1 may alter the cloud ionization fraction either by ionizing hydrogen or gaining electrons so that they change its own incident state of charge, as measured for massive ions on gas [29]. For low velocities ($\beta = v/c$) this may be important as it diminishes the ionizing power from the bare charge Z to an effective $Z_1 = Z(1 - e^{-(0.95\beta/\alpha Z^{2/3}})$ [30].

Interaction in the HI cloud may arise due to a variety of processes [28][31]. The strangelet (kinetic) energy loss per unit length due to ionization and pion production can be obtained from the stopping power $\frac{dT}{dx} = \frac{dT}{dt}\frac{1}{\beta c}$. At lower energies, ionization is the most important process, and for an incoming particle with effective charge Z_1 and velocity β the energy loss rate reads [32]

$$\frac{dT}{dt} = -1.82 \times 10^{-3} \left(\frac{n}{10^4\,{\rm cm}^{-3}}\right) [Z_1^2 \Psi(Z,\beta) + \Xi(A,\beta)] \ {\rm eV/s}, \tag{4}$$

where

$$\Psi(Z,\beta) = \frac{[1 + 0.0185 \ln(\beta)\theta(\beta - \beta_0)]}{\beta_0^3 + 2\beta^3} 2\beta^2, \tag{5}$$

and

$$\Xi(A,\beta) = 0.72 A^{0.53} \gamma^{1.28} \theta(\gamma - 1.3), \tag{6}$$

$\theta(x)$ is the Heaviside function, $\gamma^{-1} = \sqrt{1-\beta^2}$ and $\beta_0 = 0.01$ is the electron orbital velocity. This function is shown in Fig. 2 versus the strangelet kinetic energy per baryon number for an $A = 10^3$ case. Different amounts of charge $Z/A = 5\,10^{-3}$ (solid), ordinary (dotted) and CFL (dot-dashed line) strangelets and a CR proton case (dashed line) have been considered. For relativistic strangelets most of the energy is kinetic and they show an enhanced ionizing power with respect to the linear case and a change in slope around $T/A \sim 500$ MeV due to pion production for larger energies.

To estimate the hydrogen ionization rate by strangelets, we consider the number density of old NSs in our galaxy to be $n_{\rm oldNS} \sim 10^{-4}{\rm pc}^{-3}$, assuming $\sim 10^8$ NSs in a sphere of radius ~ 10 kpc. The number density of molecular clouds (MC) is $n_{MC} \sim 10^{-6}\,{\rm pc}^{-3}$, so we require on average that a fraction $f_{\rm conv} \sim 1\%$ of old NSs convert to quark stars and be sources of ionizing strangelets. The strangelet injection rate can be obtained using the production rate per source from Eq. (1), $\rm dN_A/dt\, n_{oldNS} f_{conv} \sim 2 \times 10^{39} A^{-1} yr^{-1} pc^{-3} = 2.3 \times 10^{-24} A^{-1} cm^{-3} s^{-1}$.

Taking an escape time from the MC by diffusion of the ~few GeV/A strangelet as $\Delta t \sim Z^{1/3}\, 10^2\,{\rm yr} \sim 10^{10}$ s, the strangelet number density is $n_{\rm STR} = 2.3 \times 10^{-14} Z^{1/3} A^{-1} {\rm cm}^{-3}$. This is much lower than the CR density at the low-energy break in the spectrum of Galactic CRs at ~ 1 GeV, computed from the CR flux $F_{\rm CR} \sim 10^4 {\rm m}^{-2}\,{\rm s}^{-1}\,{\rm sr}^{-1}$ or $n_{\rm CR} \sim 3\,10^{-10}\,{\rm cm}^{-3}$ [33].

The hydrogen ionization rate ζ^H *averaged over the molecular cloud*, is now estimated to be $\zeta^H = (dT^{\rm ion}/dt)\, n_{\rm STR} \langle I \rangle^{-1} n_H^{-1}$, per H atom assuming $\langle I \rangle \simeq 36$ eV per H ionization. Then $\zeta^H = Z^2 1.2 \times 10^{-25}\, s^{-1}$, per HI atom for a strangelet with $A = 10^3$.

The strangelet mean free path or average range can be calculated as $R_{\rm STR} \sim \int dT \left(\frac{dT}{dx}\right)^{-1}$. If strangelets have a net charge Z then the new range will be smaller since $R_{\rm STR} = R_{STR\,(Z=1)}/Z^2$, and therefore will provide a more efficient ionization. In Fig. 3 we show the logarithm of the strangelet range in the cloud as a function of the logarithm of the incoming kinetic energy per baryon number. We consider a proton CR case $Z = 1$, ordinary and CFL strangelets with $A = 10^2$ and $A = 10^3$ respectively, and larger strangelets with $A = 10^{13}$ and $A = 10^{18}$ with $Z/A = 10^{-7}$. As seen the cloud can effectively stop the very heavy nuggets while this is not possible for the smaller droplets even at lower kinetic energies.

Once the cloud core is populated with trails of HII (ionized hydrogen), this region contains $\rm H^+$ ions that tend to recombine radiatively as $\rm H^+ + e^- \rightarrow H + h\nu$.



FIG. 4. Logarithm of the electron column density in the cloud as a function of the logarithm of the incoming kinetic energy per baryon number. We consider a CR proton $Z = 1$, ordinary strangelet with $A = 10^2$, and large strangelets with $A = 10^9$ and $Z/A = 5 \times 10^{-4}$, $A = 10^{14}$ and $Z/A = 10^{-3}$ and $A = 10^{15}$ and $Z/A = 10^{-4}$.

The balance between ionization and recombination determines the ionized fraction. To estimate the possibility of a net ionization in the HI cloud one must compare the ionization and recombination times, $\tau_{\rm ioniz}$ and $\tau_{\rm rec}$ respectively, so that $\tau_{\rm ioniz} \lesssim \tau_{\rm rec}$. If recombination is slower, there would be a net amount of ionized HI in the cloud to cause an electron density enhancement. In addition, there might be a fraction of net charge due to the accumulation of positively charged lumps stopped by the cloud. The recombination time, $\tau_{\rm rec}$, is calculated in the *on the spot* approximation as $\tau_{\rm rec} = 1/n_e\alpha^{(2)} = 3.85 \times 10^{12} n_e^{-1} T_4^{0.8}\ s$ where $T_4 = T/10^4$ K. We estimate the average ionization time from the average number of ionization in the cloud lifetime as $\tau_{\rm ioniz} \simeq \tau_c/N_A^{\rm eff}$. Since $t_{\rm ioniz} << t_{\rm rec}$ then it is indeed possible to obtain net ionization. The ionization of interest for the ESEs occurs along the trajectory, that spans the MC, but only over a trajectory width given by the product of hydrogen recombination time and sound speed, $c_s \sim 10^6\,{\rm cm\,s^{-1}}$, namely $h \sim c_s \alpha_H^{-1} n_H^{-1} \sim 10^{15} n_4^{-1}$cm. Therefore the local ionization rate is increased with respect to a $Z = 1$ CR by $\sim Z^2 (R_{\rm C}/h)^2 \sim 10^9 Z^2\, n_4^2$, to give $\zeta^H \sim Z^2\, 10^{-16} s^{-1} n_4^2$, per HI atom. Equating $t_{rec}^{-1} = \alpha n_e$ with ζ_H yields an electron density of $n_e \sim 4 \times 10^{-4}\, Z^2 n_4^2\,{\rm cm^{-3}}$. Typical events have been argued to be explained by electron densities when the MC gas will be fully ionized in a small volume around the track of about [34] $n_e = (100 - 400)\,{\rm cm^{-3}}$, with some cases up to $n_e = 10^4\,{\rm cm^{-3}}$. Assuming an $A = 10^3$ nugget and i. e. $Z = 1 - 30$ depending of its nature, this yields trail widths of $h \sim 3.3 \times (10^{-5} - 10^{-6}) R_C \sim 3 \times 10^{13} - 3 \times 10^{12}$ cm as required by the ESE events.

The electron column density, ρ_e, can be calculated along the line of sight (los) as $\rho_e = \int_{\rm los} n_e(s) ds$, then the enhancement coming from the ionization produced by the incoming particles is estimated as $\Delta\rho_e \simeq \int_{\rm los} \Delta n_e ds$. In Fig. 4 we show the logarithm of the electron column density in the cloud as a function of the logarithm of the incoming kinetic energy per baryon number. We consider a CR proton $Z = 1$ (dashed line), ordinary strangelet with $A = 10^2$ (dotted), and large strangelets with $A = 10^9$ and $Z/A = 5 \times 10^{-4}$ (short dashed), $A = 10^{14}$ and $Z/A = 10^{-3}$ (solid) and $A = 10^{15}$ and $Z/A = 10^{-4}$ (dot-dashed). We have used the ionization produced with $N_A^{\rm eff}$ particles, the more heavy and charged the more effective are the particles at causing scintillation. These values are compatible with some quantitative models of ESE [8].

We have shown that pulsar scintillations may be caused by an ionization agent constituted by positively charged lumps of strange matter. Typical obtained over-densities in electron column densities are compatible with those of quantitative ESE models. Compact A. U. sized regions are involved in the geometry of these events. Since these quark nuggets remain to be experimentally discovered further work is needed to experimentally confirm this theoretical scenario.

M. A. P. G. would like to thank the kind hospitality of Institute d'Astrophysique in Paris where part of this work was developed and the Spanish MICINN MULTIDARK, FIS2012-30926 and FIS2011-14759-E projects.

BIBLIOGRAPHY

[1] Fiedler R. L., Dennison B., Johnston K. J., Hewish A., 1987, Nat, 326, 675

[2] Fiedler, R.; Dennison, B.; Johnston, K. J.; Waltman, E. B.; Simon, R. S. 1994ApJ...430..581

[3] Cordes, J. M., Rickett, B. J., Stinebring, D. R., Coles, W. A. 2006, ApJ, 637, 346

[4] Walker, M. A.; Melrose, D. B.; Stinebring, D. R.; Zhang, C. M. 2004, MNRAS, 354, 43

[5] Lazio, T. J. W., & Cordes, J. M. 1998, ApJ, 505, 715

[6] Goldreich P., Sridhar S., 2006, ApJ, 640, L159

[7] Walker, Mark A. 2001Ap&SS.278..149

[8] M. Walker arxiv: 0610737, published in SINS - Small Ionized and Neutral Structures in the Diffuse Interstellar Medium, ASP Conference Series, Vol. 365, eds. M. Haverkorn and W. M. Goss. San Francisco: Astronomical Society of the Pacific, 2007., p.299

[9] Revaz Y., Pfenniger D., Combes F., Bournaud F., 2009, A&A, 501, 171

[10] Walker M. A., de Bruyn A. G., Bignall H. E., 2009, MNRAS, 397, 447

[11] U. Pen, L. King , arXiv:1111.6806v1

[12] Farhi E. and Jaffe R., Phys. Rev. D 30 (1984) 2379

[13] Lu Z., Holt, R. J., Mueller, P.,Nucl. Phys. A 754 (2005) 361

[14] Perez-Garcia, A., Silk, J., Stone, J. R. , Phys. Rev. Lett. 105 (2010) 141101

[15] Perez-Garcia, M. A., Silk, J. , Physics Letters B 711, (2012), 6

[16] Perez-Garcia, M. A., Daigne, F., Silk, J. 2013, ApJ in press.

[17] Madsen, J. arXiv:astro-ph/0612784, *Strangelets in Cosmic Rays*, Invited talk at 11th Marcel Grossmann Meeting, Berlin, July 2006
[18] Shaham, N., Piran, T. arXiv:1204.1488
[19] Kotera, K. , Perez-Garcia, M. A., Silk, J., arXiv:1303.1186
[20] Spitzer, L. Tomasko, M. G. 1968, ApJ, 152 (1968) 971
[21] Glassgold, A. E., Langer, W. D., ApJ 193 (1974) 73
[22] Hotokezaka K. et al, arXiv:1212.0905v1 [astro-ph.HE]
[23] J. Schaffner-Bielich et al., Phys. Rev. C55 3038 (1997)
[24] G. Giacomelli (ANTARES collab), in press Journal of Communication and Computers, 2012.
[25] B. Christy et al (Iceube collab), Proceedings of the 30th International Cosmic ray Conference, Merida, Mexico 2007
[26] Brisken W. F. et al, ApJ, 708, (2010), 232.
[27] Madsen, J., Phys. Rev. Lett. 87 (2001) 172003
[28] Madsen, J., Phys. Rev. Lett. 85 (2000) 4687
[29] W. Liu et al, Nucl. Instr. and Meth. A 496 (2003) 198.
[30] Pierce T. E., Blann M., Phys. Rev. 173 (1968) 390, Hubert F. et al., Atom. Dat. Nucl. Dat. Tab. 46 (1990).
[31] M.S. Berger, R. L. Jaffe, PRC 35 (1987) 213
[32] Schlickeiser, R., *Cosmic Ray Astrophysics*, Astronomy and Astrophysics Library, Springer Ed., 2002 ,ISBN 978-3-540-66465-9
[33] A.Neronov, D.V.Semikoz, A.M.Taylor , Phys. Rev. Lett.108, 051105 (2012)
[34] M. Pohl, W. Reich, T. P. Krichbaum et al, arXiv:astro-ph/9503082